

## Grade 3 Learning Acceleration Guidance

Learning acceleration will ensure students have the skills they need to equitably access and practice on-grade level content.. This chart is a reference guide for teachers to help them more quickly identify the specific prerequisite and co-requisite standards necessary for every Grade 3 math standard. Students should spend the large majority of their time on the major work of the grade (■). Supporting work (■) and, where appropriate, additional work (■) can engage students in the major work of the grade.

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<b>3.OA.A.1</b> Interpret products of whole numbers, e.g., interpret $5 \times 7$ as the total number of objects in 5 groups of 7 objects each. <i>For example, describe a context in which a total number of objects can be expressed as <math>5 \times 7</math>.</i>	<b>2.OA.C.3</b> Determine whether a group of objects (up to 20) has an odd or even number of members, e.g., by pairing objects or counting them by 2s; write an equation to express an even number as a sum of two equal addends. <b>2.OA.C.4</b> Use addition to find the total number of objects arranged in rectangular arrays with up to 5 rows and up to 5 columns; write an equation to express the total as a sum of equal addends.		<b>3.OA.B.6</b> Understand division as an unknown-factor problem. <i>For example, find <math>32 \div 8</math> by finding the number that makes 32 when multiplied by 8.</i>
<b>3.OA.A.2</b> Interpret whole-number quotients of whole numbers, e.g., interpret $56 \div 8$ as the number of objects in each share when 56 objects are partitioned equally into 8 shares, or as a number of shares when 56 objects are partitioned into equal shares of 8 objects each. <i>For example, describe a context in which a number of shares or a number of groups can be expressed as <math>56 \div 8</math>.</i>		<b>3.OA.A.1</b> Interpret products of whole numbers, e.g., interpret $5 \times 7$ as the total number of objects in 5 groups of 7 objects each. <i>For example, describe a context in which a total number of objects can be expressed as <math>5 \times 7</math>.</i>	<b>3.OA.B.6</b> Understand division as an unknown-factor problem. <i>For example, find <math>32 \div 8</math> by finding the number that makes 32 when multiplied by 8.</i>

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<p><b>3.OA.A.3</b> Use multiplication and division within 100 to solve word problems in situations involving equal groups, arrays, and measurement quantities, e.g., by using drawings and equations with a symbol for the unknown number to represent the problem.</p>		<p><b>3.OA.A.1</b> Interpret products of whole numbers, e.g., interpret <math>5 \times 7</math> as the total number of objects in 5 groups of 7 objects each. <i>For example, describe a context in which a total number of objects can be expressed as <math>5 \times 7</math>.</i></p> <p><b>3.OA.A.2</b> Interpret whole-number quotients of whole numbers, e.g., interpret <math>56 \div 8</math> as the number of objects in each share when 56 objects are partitioned equally into 8 shares, or as a number of shares when 56 objects are partitioned into equal shares of 8 objects each. <i>For example, describe a context in which a number of shares or a number of groups can be expressed as <math>56 \div 8</math>.</i></p>	<p><b>3.OA.A.4</b> Determine the unknown whole number in a multiplication or division equation relating three whole numbers. <i>For example, determine the unknown number that makes the equation true in each of the equations <math>8 \times ? = 48</math>, <math>5 = \_ \div 3</math>, <math>6 \times 6 = ?</math></i></p> <p><b>3.OA.B.6</b> Understand division as an unknown-factor problem. <i>For example, find <math>32 \div 8</math> by finding the number that makes 32 when multiplied by 8.</i></p>
<p><b>3.OA.A.4</b> Determine the unknown whole number in a multiplication or division equation relating three whole numbers. <i>For example, determine the unknown number that makes the equation true in each of the equations <math>8 \times ? = 48</math>, <math>5 = \_ \div 3</math>, <math>6 \times 6 = ?</math></i></p>			<p><b>3.OA.A.3</b> Use multiplication and division within 100 to solve word problems in situations involving equal groups, arrays, and measurement quantities, e.g., by using drawings and equations with a symbol for the unknown number to represent the problem.</p> <p><b>3.OA.C.7</b> Fluently multiply and divide within 100, using strategies such as the relationship between multiplication and division (e.g., knowing that <math>8 \times 5 = 40</math>, one knows <math>40 \div 5 = 8</math>) or properties of operations. By the end of Grade 3, know from memory all products of two one-digit numbers.</p>
<p><b>3.OA.B.5</b> Apply properties of operations as strategies to multiply and divide.<sup>2</sup> <i>Examples: If <math>6 \times 4 = 24</math> is known, then <math>4 \times 6 = 24</math> is also known. (Commutative property of multiplication.) <math>3 \times 5 \times 2</math> can be found by <math>3 \times 5 = 15</math>, then <math>15 \times 2 = 30</math>, or by <math>5 \times 2 = 10</math>, then <math>3 \times 10 = 30</math>. (Associative property of multiplication.) Knowing that <math>8 \times 5 = 40</math> and <math>8 \times 2 = 16</math>, one can find <math>8 \times 7</math> as <math>8 \times (5 + 2) = (8 \times 5) + (8 \times 2) = 40 + 16 = 56</math>. (Distributive property.)</i></p>		<p><b>3.OA.A.1</b> Interpret products of whole numbers, e.g., interpret <math>5 \times 7</math> as the total number of objects in 5 groups of 7 objects each. <i>For example, describe a context in which a total number of objects can be expressed as <math>5 \times 7</math>.</i></p> <p><b>3.OA.A.2</b> Interpret whole-number quotients of whole numbers, e.g., interpret <math>56 \div 8</math> as the number of objects in each share when 56 objects are partitioned equally into 8 shares, or as a number of shares when 56 objects are partitioned into equal shares of 8 objects each. <i>For example, describe a context in which a number of shares or a number of groups can be expressed as <math>56 \div 8</math>.</i></p>	

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<p><b>3.OA.B.6</b> Understand division as an unknown-factor problem. <i>For example, find <math>32 \div 8</math> by finding the number that makes 32 when multiplied by 8.</i></p>			<p><b>3.OA.A.1</b> Interpret products of whole numbers, e.g., interpret <math>5 \times 7</math> as the total number of objects in 5 groups of 7 objects each. <i>For example, describe a context in which a total number of objects can be expressed as <math>5 \times 7</math>.</i></p> <p><b>3.OA.A.2</b> Interpret whole-number quotients of whole numbers, e.g., interpret <math>56 \div 8</math> as the number of objects in each share when 56 objects are partitioned equally into 8 shares, or as a number of shares when 56 objects are partitioned into equal shares of 8 objects each. <i>For example, describe a context in which a number of shares or a number of groups can be expressed as <math>56 \div 8</math>.</i></p> <p><b>3.OA.A.3</b> Use multiplication and division within 100 to solve word problems in situations involving equal groups, arrays, and measurement quantities, e.g., by using drawings and equations with a symbol for the unknown number to represent the problem.</p>
<p><b>3.OA.C.7</b> Fluently multiply and divide within 100, using strategies such as the relationship between multiplication and division (e.g., knowing that <math>8 \times 5 = 40</math>, one knows <math>40 \div 5 = 8</math>) or properties of operations. By the end of Grade 3, know from memory all products of two one-digit numbers.</p>		<p><b>3.OA.B.5</b> Apply properties of operations as strategies to multiply and divide.<sup>2</sup> <i>Examples: If <math>6 \times 4 = 24</math> is known, then <math>4 \times 6 = 24</math> is also known. (Commutative property of multiplication.) <math>3 \times 5 \times 2</math> can be found by <math>3 \times 5 = 15</math>, then <math>15 \times 2 = 30</math>, or by <math>5 \times 2 = 10</math>, then <math>3 \times 10 = 30</math>. (Associative property of multiplication.) Knowing that <math>8 \times 5 = 40</math> and <math>8 \times 2 = 16</math>, one can find <math>8 \times 7</math> as <math>8 \times (5 + 2) = (8 \times 5) + (8 \times 2) = 40 + 16 = 56</math>. (Distributive property.)</i></p> <p><b>3.OA.B.6</b> Understand division as an unknown-factor problem. <i>For example, find <math>32 \div 8</math> by finding the number that makes 32 when multiplied by 8.</i></p>	<p><b>3.OA.A.4</b> Determine the unknown whole number in a multiplication or division equation relating three whole numbers. <i>For example, determine the unknown number that makes the equation true in each of the equations <math>8 \times ? = 48</math>, <math>5 = \_ \div 3</math>, <math>6 \times 6 = ?</math></i></p> <p><b>3.OA.D.8</b> Solve two-step word problems using the four operations. Represent these problems using equations with a letter standing for the unknown quantity. Assess the reasonableness of answers using mental computation and estimation strategies including rounding.</p>

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<p><b>3.OA.D.8</b> Solve two-step word problems using the four operations. Represent these problems using equations with a letter standing for the unknown quantity. Assess the reasonableness of answers using mental computation and estimation strategies including rounding.</p>	<p><b>2.OA.A.1</b> Use addition and subtraction within 100 to solve one- and two-step word problems involving situations of adding to, taking from, putting together, taking apart, and comparing, with unknowns in all positions, e.g., by using drawings and equations with a symbol for the unknown number to represent the problem.</p>	<p><b>3.OA.A.3</b> Use multiplication and division within 100 to solve word problems in situations involving equal groups, arrays, and measurement quantities, e.g., by using drawings and equations with a symbol for the unknown number to represent the problem.</p>	<p><b>3.OA.C.7</b> Fluently multiply and divide within 100, using strategies such as the relationship between multiplication and division (e.g., knowing that <math>8 \times 5 = 40</math>, one knows <math>40 \div 5 = 8</math>) or properties of operations. By the end of Grade 3, know from memory all products of two one-digit numbers.</p> <p><b>3.MD.A.2</b> Measure and estimate liquid volumes and masses of objects using standard units of grams (g), kilograms (kg), and liters (l). Add, subtract, multiply, or divide to solve one-step word problems involving masses or volumes that are given in the same units, e.g., by using drawings (such as a beaker with a measurement scale) to represent the problem.</p> <p><b>3.MD.B.3</b> Draw a scaled picture graph and a scaled bar graph to represent a data set with several categories. Solve one- and two-step “how many more” and “how many less” problems using information presented in scaled bar graphs. For example, draw a bar graph in which each square in the bar graph might represent 5 pets.</p> <p><b>3.MD.D.8</b> Solve real-world and mathematical problems involving perimeters of polygons, including finding the perimeter given the side lengths, finding an unknown side length, and exhibiting rectangles with the same perimeter and different areas or with the same area and different perimeters.</p>
<p><b>3.OA.D.9</b> Identify arithmetic patterns (including patterns in the addition table or multiplication table), and explain them using properties of operations. For example, observe that 4 times a number is always even, and explain why 4 times a number can be decomposed into two equal addends.</p>	<p><b>2.OA.C.3</b> Determine whether a group of objects (up to 20) has an odd or even number of members, e.g., by pairing objects or counting them by 2s; write an equation to express an even number as a sum of two equal addends.</p>	<p><b>3.OA.B.5</b> Apply properties of operations as strategies to multiply and divide.<sup>2</sup> <i>Examples: If <math>6 \times 4 = 24</math> is known, then <math>4 \times 6 = 24</math> is also known. (Commutative property of multiplication.) <math>3 \times 5 \times 2</math> can be found by <math>3 \times 5 = 15</math>, then <math>15 \times 2 = 30</math>, or by <math>5 \times 2 = 10</math>, then <math>3 \times 10 = 30</math>. (Associative property of multiplication.) Knowing that <math>8 \times 5 = 40</math> and <math>8 \times 2 = 16</math>, one can find <math>8 \times 7</math> as <math>8 \times (5 + 2) = (8 \times 5) + (8 \times 2) = 40 + 16 = 56</math>. (Distributive property.)</i></p>	

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<p><b>3.NBT.A.1</b> Use place value understanding to round whole numbers to the nearest 10 or 100.</p>	<p><b>2.NBT.A.1</b> 1. Understand that the three digits of a three-digit number represent amounts of hundreds, tens, and ones; e.g., 706 equals 7 hundreds, 0 tens, and 6 ones. Understand the following as special cases: a. 100 can be thought of as a bundle of ten tens—called a “hundred.” b. The numbers 100, 200, 300, 400, 500, 600, 700, 800, 900 refer to one, two, three, four, five, six, seven, eight, or nine hundreds (and 0 tens and 0 ones).</p>		
<p><b>3.NBT.A.2</b> Fluently add and subtract within 1000 using strategies and algorithms based on place value, properties of operations, and/or the relationship between addition and subtraction.</p>	<p><b>2.NBT.B.7</b> Add and subtract within 1000, using concrete models or drawings and strategies based on place value, properties of operations, and/or the relationship between addition and subtraction; justify the reasoning used with a written explanation. Understand that in adding or subtracting three-digit numbers, one adds or subtracts hundreds and hundreds, tens and tens, ones and ones; and sometimes it is necessary to compose or decompose tens or hundreds. <b>2.NBT.B.8</b> Mentally add 10 or 100 to a given number 100–900, and mentally subtract 10 or 100 from a given number 100–900.</p>		
<p><b>3.NBT.A.3</b> Multiply one-digit whole numbers by multiples of 10 in the range 10–90 (e.g., <math>9 \times 80</math>, <math>5 \times 60</math>) using strategies based on place value and properties of operations.</p>	<p><b>2.NBT.A.1</b> 1. Understand that the three digits of a three-digit number represent amounts of hundreds, tens, and ones; e.g., 706 equals 7 hundreds, 0 tens, and 6 ones. Understand the following as special cases: a. 100 can be thought of as a bundle of ten tens—called a “hundred.” b. The numbers 100, 200, 300, 400, 500, 600, 700, 800, 900 refer to one, two, three, four, five, six, seven, eight, or nine hundreds (and 0 tens and 0 ones).</p>	<p><b>3.OA.B.5</b> Apply properties of operations as strategies to multiply and divide.<sup>2</sup> <i>Examples: If <math>6 \times 4 = 24</math> is known, then <math>4 \times 6 = 24</math> is also known. (Commutative property of multiplication.) <math>3 \times 5 \times 2</math> can be found by <math>3 \times 5 = 15</math>, then <math>15 \times 2 = 30</math>, or by <math>5 \times 2 = 10</math>, then <math>3 \times 10 = 30</math>. (Associative property of multiplication.) Knowing that <math>8 \times 5 = 40</math> and <math>8 \times 2 = 16</math>, one can find <math>8 \times 7</math> as <math>8 \times (5 + 2) = (8 \times 5) + (8 \times 2) = 40 + 16 = 56</math>. (Distributive property.)</i></p>	

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<p><b>3.NF.A.1</b></p> <p>Understand a fraction <math>1/b</math>, with denominators 2, 3, 4, 6, and 8, as the quantity formed by 1 part when a whole is partitioned into <math>b</math> equal parts; understand a fraction <math>a/b</math> as the quantity formed by <math>a</math> parts of size <math>1/b</math>.</p>	<p><b>2.MD.A.2</b></p> <p>Measure the length of an object twice, using length units of different lengths for the two measurements; describe how the two measurements relate to the size of the unit chosen.</p> <p><b>2.G.A.3</b></p> <p>Partition circles and rectangles into two, three, or four equal shares, describe the shares using the words halves, thirds, half of, a third of, etc., and describe the whole as two halves, three thirds, four fourths. Recognize that equal shares of identical wholes need not have the same shape.</p>		<p><b>3.NF.A.2</b></p> <p>Understand a fraction with denominators 2, 3, 4, 6, and 8 as a number on the number line; represent fractions on a number line diagram.</p> <p><b>3.MD.A.2</b></p> <p>Measure and estimate liquid volumes and masses of objects using standard units of grams (g), kilograms (kg), and liters (l). Add, subtract, multiply, or divide to solve one-step word problems involving masses or volumes that are given in the same units, e.g., by using drawings (such as a beaker with a measurement scale) to represent the problem.</p>
<p><b>3.NF.A.2</b></p> <p>Understand a fraction with denominators 2, 3, 4, 6, and 8 as a number on a number line diagram.</p> <p>a. Represent a fraction <math>1/b</math> on a number line diagram by defining the interval from 0 to 1 as the whole and partitioning it into <math>b</math> equal parts. Recognize that each part has size <math>1/b</math> and that the endpoint of the part based at 0 locates the number <math>1/b</math> on the number line.</p> <p>b. Represent a fraction <math>a/b</math> on a number line diagram by marking off <math>a</math> lengths <math>1/b</math> from 0. Recognize that the resulting interval has size <math>a/b</math> and that its endpoint locates the number <math>a/b</math> on the number line.</p>	<p><b>2.MD.B.6</b></p> <p>Represent whole numbers as lengths from 0 on a number line diagram with equally spaced points corresponding to the numbers 0, 1, 2, ..., and represent whole-number sums and differences within 100 on a number line diagram.</p>		<p><b>3.NF.A.1</b></p> <p>Understand a fraction <math>1/b</math>, with denominators 2, 3, 4, 6, and 8, as the quantity formed by 1 part when a whole is partitioned into <math>b</math> equal parts; understand a fraction <math>a/b</math> as the quantity formed by <math>a</math> parts of size <math>1/b</math>.</p> <p><b>3.MD.B.4</b></p> <p>Generate measurement data by measuring lengths using rulers marked with halves and fourths of an inch. Show the data by making a line plot, where the horizontal scale is marked off in appropriate units— whole numbers, halves, or quarters.</p>

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<p><b>3.NF.A.3</b>            Explain equivalence of fractions with denominators 2, 3, 4, 6, and 8 in special cases, and compare fractions by reasoning about their size.</p> <ul style="list-style-type: none"> <li>a. Understand two fractions as equivalent (equal) if they are the same size, or the same point on a number line.</li> <li>b. Recognize and generate simple equivalent fractions, e.g., <math>1/2 = 2/4</math>, <math>4/6 = 2/3</math>. Explain why the fractions are equivalent, e.g., by using a visual fraction model.</li> <li>c. Express whole numbers as fractions, and recognize fractions that are equivalent to whole numbers. Examples: Express 3 in the form <math>3 = 3/1</math>; recognize that <math>6/1 = 6</math>; locate <math>4/4</math> and 1 at the same point of a number line diagram.</li> <li>d. Compare two fractions with the same numerator or the same denominator by reasoning about their size. Recognize that comparisons are valid only when the two fractions refer to the same whole. Record the results of comparisons with the symbols <math>&gt;</math>, <math>=</math>, or <math>&lt;</math>, and justify the conclusions, e.g., by using a visual fraction model.</li> </ul>		<p><b>3.NF.A.1</b>            Understand a fraction <math>1/b</math>, with denominators 2, 3, 4, 6, and 8, as the quantity formed by 1 part when a whole is partitioned into <math>b</math> equal parts; understand a fraction <math>a/b</math> as the quantity formed by <math>a</math> parts of size <math>1/b</math>.</p> <p><b>3.NF.A.2</b>            Understand a fraction with denominators 2, 3, 4, 6, and 8 as a number on the number line; represent fractions on a number line diagram.</p> <ul style="list-style-type: none"> <li>a. Represent a fraction <math>1/b</math> on a number line diagram by defining the interval from 0 to 1 as the whole and partitioning it into <math>b</math> equal parts. Recognize that each part has size <math>1/b</math> and that the endpoint of the part based at 0 locates the number <math>1/b</math> on the number line.</li> <li>b. Represent a fraction <math>a/b</math> on a number line diagram by marking off <math>a</math> lengths <math>1/b</math> from 0. Recognize that the resulting interval has size <math>a/b</math> and that its endpoint locates the number <math>a/b</math> on the number line.</li> </ul>	
<p><b>3.MD.A.1</b>            Understand time to the nearest minute.</p> <ul style="list-style-type: none"> <li>a. Tell and write time to the nearest minute and measure time intervals in minutes, within 60 minutes, on an analog and digital clock.</li> <li>b. Calculate elapsed time greater than 60 minutes to the nearest quarter and half hour on a number line diagram.</li> <li>c. Solve word problems involving addition and subtraction of time intervals in minutes, e.g., by representing the problem on a number line diagram.</li> </ul>			

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<p><b>3.MD.A.2</b> Measure and estimate liquid volumes and masses of objects using standard units of grams (g), kilograms (kg), and liters (l). Add, subtract, multiply, or divide to solve one-step word problems involving masses or volumes that are given in the same units, e.g., by using drawings (such as a beaker with a measurement scale) to represent the problem.</p>	<p><b>2.MD.A.1</b> Measure the length of an object by selecting and using appropriate tools such as rulers, yardsticks, meter sticks, and measuring tapes.</p>		<p><b>3.NF.A.1</b> Understand a fraction <math>1/b</math>, with denominators 2, 3, 4, 6, and 8, as the quantity formed by 1 part when a whole is partitioned into <math>b</math> equal parts; understand a fraction <math>a/b</math> as the quantity formed by <math>a</math> parts of size <math>1/b</math>.</p> <p><b>3.OA.D.8</b> Solve two-step word problems using the four operations. Represent these problems using equations with a letter standing for the unknown quantity. Assess the reasonableness of answers using mental computation and estimation strategies including rounding.</p>
<p><b>3.MD.B.3</b> Draw a scaled picture graph and a scaled bar graph to represent a data set with several categories. Solve one- and two-step “how many more” and “how many less” problems using information presented in scaled bar graphs. For example, draw a bar graph in which each square in the bar graph might represent 5 pets.</p>			<p><b>3.OA.D.8</b> Solve two-step word problems using the four operations. Represent these problems using equations with a letter standing for the unknown quantity. Assess the reasonableness of answers using mental computation and estimation strategies including rounding.</p>
<p><b>3.MD.B.4</b> Generate measurement data by measuring lengths using rulers marked with halves and fourths of an inch. Show the data by making a line plot, where the horizontal scale is marked off in appropriate units— whole numbers, halves, or quarters.</p>			<p><b>3.NF.A.2</b> Understand a fraction with denominators 2, 3, 4, 6, and 8 as a number on the number line; represent fractions on a number line diagram.</p> <ol style="list-style-type: none"> <li>Represent a fraction <math>1/b</math> on a number line diagram by defining the interval from 0 to 1 as the whole and partitioning it into <math>b</math> equal parts. Recognize that each part has size <math>1/b</math> and that the endpoint of the part based at 0 locates the number <math>1/b</math> on the number line.</li> <li>Represent a fraction <math>a/b</math> on a number line diagram by marking off <math>a</math> lengths <math>1/b</math> from 0. Recognize that the resulting interval has size <math>a/b</math> and that its endpoint locates the number <math>a/b</math> on the number line.</li> </ol>

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<p><b>3.MD.C.5</b> Recognize area as an attribute of plane figures and understand concepts of area measurement.</p> <p>a. A square with side length 1 unit, called “a unit square,” is said to have “one square unit” of area, and can be used to measure area.</p> <p>b. A plane figure which can be covered without gaps or overlaps by <math>n</math> unit squares is said to have an area of <math>n</math> square units.</p>	<p><b>1.G.A.2</b> Compose two-dimensional shapes (rectangles, squares, trapezoids, triangles, half-circles, and quarter-circles) and three-dimensional shapes (cubes, right rectangular prisms, right circular cones, and right circular cylinders) to create a composite shape, and compose new shapes from the composite shape.</p> <p><b>2.MD.A.1</b> Measure the length of an object by selecting and using appropriate tools such as rulers, yardsticks, meter sticks, and measuring tapes.</p>		
<p><b>3.MD.C.6</b> Measure areas by counting unit squares (square cm, square m, square in, square ft, and improvised units).</p>	<p><b>2.G.A.2</b> Partition a rectangle into rows and columns of same-size squares and count to find the total number of them.</p>	<p><b>3.MD.C.5</b> Recognize area as an attribute of plane figures and understand concepts of area measurement.</p> <p>a. A square with side length 1 unit, called “a unit square,” is said to have “one square unit” of area, and can be used to measure area.</p> <p>b. A plane figure which can be covered without gaps or overlaps by <math>n</math> unit squares is said to have an area of <math>n</math> square units.</p>	
<p><b>3.MD.C.7</b> Relate area to the operations of multiplication and addition.</p> <p>a. Find the area of a rectangle with whole-number side lengths by tiling it, and show that the area is the same as would be found by multiplying the side lengths.</p> <p>b. Multiply side lengths to find areas of rectangles with whole- number side lengths in the context of solving real-world and mathematical problems, and represent whole-number products as rectangular areas in mathematical reasoning.</p> <p>c. Use tiling to show in a concrete case that the area of a rectangle with whole-number side lengths <math>a</math> and <math>b + c</math> is the sum of <math>a + b</math> and <math>a + c</math>. Use area models to represent the distributive property in mathematical reasoning.</p>		<p><b>3.MD.C.5</b> Recognize area as an attribute of plane figures and understand concepts of area measurement.</p> <p>a. A square with side length 1 unit, called “a unit square,” is said to have “one square unit” of area, and can be used to measure area.</p> <p>b. A plane figure which can be covered without gaps or overlaps by <math>n</math> unit squares is said to have an area of <math>n</math> square units.</p> <p><b>3.MD.C.6</b> Measure areas by counting unit squares (square cm, square m, square in, square ft, and improvised units).</p>	<p><b>3.OA.B.5</b> Apply properties of operations as strategies to multiply and divide.<sup>2</sup> <i>Examples: If <math>6 \times 4 = 24</math> is known, then <math>4 \times 6 = 24</math> is also known. (Commutative property of multiplication.) <math>3 \times 5 \times 2</math> can be found by <math>3 \times 5 = 15</math>, then <math>15 \times 2 = 30</math>, or by <math>5 \times 2 = 10</math>, then <math>3 \times 10 = 30</math>. (Associative property of multiplication.) Knowing that <math>8 \times 5 = 40</math> and <math>8 \times 2 = 16</math>, one can find <math>8 \times 7</math> as <math>8 \times (5 + 2) = (8 \times 5) + (8 \times 2) = 40 + 16 = 56</math>. (Distributive property.)</i></p> <p><b>3.OA.D.8</b> Solve two-step word problems using the four operations. Represent these problems using equations with a letter standing for the unknown quantity. Assess the reasonableness of answers using mental computation and estimation strategies including rounding.</p>

3 <sup>rd</sup> Grade Standard	Previous Grade(s) Standards	3 <sup>rd</sup> Grade Standards Taught in Advance	3 <sup>rd</sup> Grade Standards Taught Concurrently
<p><b>3.MD.D.8</b> Solve real-world and mathematical problems involving perimeters of polygons, including finding the perimeter given the side lengths, finding an unknown side length, and exhibiting rectangles with the same perimeter and different areas or with the same area and different perimeters.</p>		<p><b>3.MD.C.5</b> Recognize area as an attribute of plane figures and understand concepts of area measurement.</p> <ol style="list-style-type: none"> <li>A square with side length 1 unit, called “a unit square,” is said to have “one square unit” of area, and can be used to measure area.</li> <li>A plane figure which can be covered without gaps or overlaps by <math>n</math> unit squares is said to have an area of <math>n</math> square units.</li> </ol>	<p><b>3.OA.D.8</b> Solve two-step word problems using the four operations. Represent these problems using equations with a letter standing for the unknown quantity. Assess the reasonableness of answers using mental computation and estimation strategies including rounding.</p>
<p><b>3.MD.E.9</b> Solve word problems involving pennies, nickels, dimes, quarters, and bills greater than one dollar, using the dollar and cent symbols appropriately.</p>	<p><b>2.MD.C.8</b> Solve word problems involving dollar bills, quarters, dimes, nickels, and pennies, using \$ and ¢ symbols appropriately. Example: If you have 2 dimes and 3 pennies, how many cents do you have?</p>		
<p><b>3.G.A.1</b> Understand that shapes in different categories (e.g., rhombuses, rectangles, and others) may share attributes (e.g., having four sides), and that the shared attributes can define a larger category (e.g., quadrilaterals). Recognize rhombuses, rectangles, and squares as examples of quadrilaterals, and draw examples of quadrilaterals that do not belong to any of these subcategories.</p>	<p><b>2.G.A.1</b> Recognize and draw shapes having specified attributes, such as a given number of angles or a given number of equal faces. Identify triangles, quadrilaterals, pentagons, hexagons, and cubes.</p>		
<p><b>3.G.A.2</b> Partition shapes into parts with equal areas. Express the area of each part as a unit fraction of the whole. For example, partition a shape into 4 parts with equal area, and describe the area of each part as <math>\frac{1}{4}</math> of the area of the shape.</p>		<p><b>3.NF.A.1</b> Understand a fraction <math>\frac{1}{b}</math>, with denominators 2, 3, 4, 6, and 8, as the quantity formed by 1 part when a whole is partitioned into <math>b</math> equal parts; understand a fraction <math>\frac{a}{b}</math> as the quantity formed by <math>a</math> parts of size <math>\frac{1}{b}</math>.</p>	