



## Grade 2 Learning Acceleration Guidance

Learning acceleration will ensure students have the skills they need to equitably access and practice on-grade level content. This chart is a reference guide for teachers to help them more quickly identify the specific prerequisite and co-requisite standards necessary for every Grade 2 math standard. Students should spend the large majority of their time on the major work of the grade ( $\blacksquare$ ). Supporting work ( $\blacksquare$ ) and, where appropriate, additional work ( $\blacksquare$ ) can engage students in the major work of the grade.

in the major work of the grade.			
2 <sup>nd</sup> Grade Standard	Previous Grade(s) Standards	2 <sup>nd</sup> Grade Standards Taught in Advance	2 <sup>nd</sup> Grade Standards Taught Concurrently
2.OA.A.1 Use addition and subtraction within 100 to solve one- and two-step word problems involving situations of adding to, taking from, putting together, taking apart, and comparing, with unknowns in all positions, e.g., by using drawings and equations with a symbol for the unknown number to represent the problem.	<ul> <li>1.OA.A.1 Use addition and subtraction within 20 to solve word problems involving situations of adding to, taking from, putting together, taking apart, and comparing, with unknowns in all positions, e.g., by using objects, drawings, and equations with a symbol for the unknown number to represent the problem. </li> <li>1.NBT.C.4 Add within 100, including adding a two-digit number and a one-digit number, and adding a two-digit number and a multiple of 10. <ul> <li>a. Use concrete models or drawings and strategies based on place value, properties of operations, and/or the relationship between addition and subtraction; relate the strategy to a number sentence; justify the reasoning used with a written explanation.</li> <li>b. Understand that in adding two-digit numbers, one adds tens and tens, ones and ones; and sometimes it is necessary to compose a ten.</li> </ul> </li> <li>1.NBT.C.5 Given a two-digit number, mentally find 10 more or 10 less than the number, without having to count; explain the reasoning used. 1.NBT.C.6 Subtract multiples of 10 in the range 10-90 (positive or zero differences), using concrete models or drawings and strategies based on place value, properties of operations, and/or the relationship between addition and subtraction; relate the strategy to a written method and explain the reasoning used.</li></ul>		<ul> <li>2.NBT.B.5</li> <li>Fluently add and subtract within 100 using strategies based on place value, properties of operations, and/or the relationship between addition and subtraction.</li> <li>2.MD.B.5</li> <li>Use addition and subtraction within 100 to solve word problems involving lengths that are given in the same units, e.g., by using drawings (such as drawings of rulers) and equations with a symbol for the unknown number to represent the problem.</li> <li>2.MD.C.8</li> <li>Solve word problems involving dollar bills, quarters, dimes, nickels, and pennies, using \$ and ¢ symbols appropriately. <i>Example: If you have 2 dimes and 3 pennies, how many cents do you have?</i></li> <li>2.MD.D.10</li> <li>Draw a picture graph and a bar graph (with single-unit scale) to represent a data set with up to four categories. Solve simple put- together, take-apart, and compare problems using information presented in a bar graph.</li> </ul>

2 <sup>nd</sup> Grade Standard	Previous Grade(s) Standards	2 <sup>nd</sup> Grade Standards Taught in Advance	2 <sup>nd</sup> Grade Standards Taught Concurrently
2.0A.B.2 Fluently add and subtract within 20 using mental strategies.2 By end of Grade 2, know from memory all sums of two one-digit numbers.	<b>1.OA.C.6</b> Add and subtract within 20, demonstrating fluency for addition and subtraction within 10. Use strategies such as counting on; making ten (e.g., $8 + 6 = 8 + 2 + 4 = 10 + 4 = 14$ ); decomposing a number leading to a ten (e.g., $13 - 4 = 13 - 3 - 1 = 10 - 1 = 9$ ); using the relationship between addition and subtraction (e.g., knowing that $8 + 4 = 12$ , one knows $12 - 8 = 4$ ); and creating equivalent but easier or known sums (e.g., adding $6 + 7$ by creating the known equivalent $6 + 6 + 1 = 12 + 4$		
2.OA.C.3 Determine whether a group of objects (up to 20) has an odd or even number of members, e.g., by pairing objects or counting them by 2s; write an equation to express an even number as a sum of two equal addends.	1 = 13). <b>1.OA.D.7</b> Understand the meaning of the equal sign, and determine if equations involving addition and subtraction are true or false. For example, which of the following equations are true and which are false? $6 = 6$ , $7 = 8 - 1$ , $5 + 2$ = 2 + 5, $4 + 1 = 5 + 2$ .		2.NBT.A.2 Count within 1000; skip-count by 5s, 10s, and 100s.
2.OA.C.4 Use addition to find the total number of objects arranged in rectangular arrays with up to 5 rows and up to 5 columns; write an equation to express the total as a sum of equal addends.	<b>1.OA.D.7</b> Understand the meaning of the equal sign, and determine if equations involving addition and subtraction are true or false. For example, which of the following equations are true and which are false? $6 = 6$ , $7 = 8 - 1$ , $5 + 2$ = 2 + 5, $4 + 1 = 5 + 2$ .		
<ul> <li>2.NBT.A.1</li> <li>Understand that the three digits of a three-digit number represent amounts of hundreds, tens, and ones; e.g., 706 equals 7 hundreds, 0 tens, and 6 ones. Understand the following as special cases: <ul> <li>a. 100 can be thought of as a bundle of ten tens — called a "hundred."</li> <li>b. The numbers 100, 200, 300, 400, 500, 600, 700, 800, 900 refer to one, two, three, four, five, six, seven, eight, or nine hundreds (and 0 tens and 0 ones).</li> </ul> </li> </ul>	<ul> <li>1.NBT.B.2</li> <li>Understand that the two digits of a two-digit number represent amounts of tens and ones.</li> <li>Understand the following as special cases: <ul> <li>a. 10 can be thought of as a bundle of ten ones — called a "ten."</li> <li>b. The numbers from 11 to 19 are composed of a ten and one, two, three, four, five, six, seven, eight, or nine ones.</li> </ul> </li> <li>c. The numbers 10, 20, 30, 40, 50, 60, 70, 80, 90 refer to one, two, three, four, five, six, seven, eight, or nine tens (and 0 ones).</li> </ul>	2.NBT.A.2 Count within 1000; skip-count by 5s, 10s, and 100s.	
2.NBT.A.2 Count within 1000; skip-count by 5s, 10s, and 100s.			2.OA.C.3 Determine whether a group of objects (up to 20) has an odd or even number of members, e.g., by pairing objects or counting them by 2s; write an equation to express an even number as a sum of two equal addends.

2 <sup>nd</sup> Grade Standard	Previous Grade(s) Standards	2 <sup>nd</sup> Grade Standards Taught in Advance	2 <sup>nd</sup> Grade Standards Taught Concurrently
2.NBT.A.3 Read and write numbers to 1000 using base- ten numerals, number names, and expanded form.		<ul> <li>2.NBT.A.1</li> <li>Understand that the three digits of a three-digit number represent amounts of hundreds, tens, and ones; e.g., 706 equals 7 hundreds, 0 tens, and 6 ones. Understand the following as special cases: <ul> <li>a. 100 can be thought of as a bundle of ten tens — called a "hundred."</li> <li>b. The numbers 100, 200, 300, 400, 500, 600, 700, 800, 900 refer to one, two, three, four, five, six, seven, eight, or nine hundreds (and 0 tens and 0 ones).</li> </ul> </li> </ul>	
2.NBT.A.4 Compare two three-digit numbers based on meanings of the hundreds, tens, and ones digits, using >, =, and < symbols to record the results of comparisons.		<ul> <li>2.NBT.A.1 Understand that the three digits of a three-digit number represent amounts of hundreds, tens, and ones; e.g., 706 equals 7 hundreds, 0 tens, and 6 ones. Understand the following as special cases: <ul> <li>a. 100 can be thought of as a bundle of ten tens — called a "hundred."</li> <li>b. The numbers 100, 200, 300, 400, 500, 600, 700, 800, 900 refer to one, two, three, four, five, six, seven, eight, or nine hundreds (and 0 tens and 0 ones).</li> </ul></li></ul>	

2 <sup>nd</sup> Grade Standard	Previous Grade(s) Standards	2 <sup>nd</sup> Grade Standards Taught in Advance	2 <sup>nd</sup> Grade Standards Taught Concurrently
2.NBT.B.5	1.NBT.C.4	2.OA.B.2	2.OA.A.1
Fluently add and subtract within 100 using	Add within 100, including adding a two-digit	Fluently add and subtract within 20 using	Use addition and subtraction within 100 to solve
strategies based on place value, properties of	number and a one-digit number, and adding a	mental strategies.2 By end of Grade 2, know	one- and two-step word problems involving
operations, and/or the relationship between	two-digit number and a multiple of 10.	from memory all sums of two one-digit	situations of adding to, taking from, putting
addition and subtraction.	<ul> <li>Use concrete models or drawings and strategies based on place value,</li> </ul>	numbers.	together, taking apart, and comparing, with
	properties of operations, and/or the		unknowns in all positions, e.g., by using drawings and equations with a symbol for the
	relationship between addition and		unknown number to represent the problem.
	subtraction; relate the strategy to a		unknown number to represent the problem.
	number sentence; justify the reasoning		
	used with a written explanation.		
	b. Understand that in adding two-digit		
	numbers, one adds tens and tens, ones		
	and ones; and sometimes it is necessary		
	to compose a ten.		
	1.NBT.C.5		
	Given a two-digit number, mentally find 10		
	more or 10 less than the number, without		
	having to count; explain the reasoning used.		
	1.NBT.C.6		
	Subtract multiples of 10 in the range 10-90		
	from multiples of 10 in the range 10-90		
	(positive or zero differences), using concrete		
	models or drawings and strategies based on place value, properties of operations, and/or		
	the relationship between addition and		
	subtraction; relate the strategy to a written		
	method and explain the reasoning used.		

2 <sup>nd</sup> Grade Standard	Previous Grade(s) Standards	2 <sup>nd</sup> Grade Standards Taught in Advance	2 <sup>nd</sup> Grade Standards Taught Concurrently
2.NBT.B.6		2.NBT.A.1	
Add up to four two-digit numbers using		Understand that the three digits of a three-	
strategies based on place value and		digit number represent amounts of hundreds,	
properties of operations.		tens, and ones; e.g., 706 equals 7 hundreds, 0	
		tens, and 6 ones. Understand the following as	
		special cases:	
		a. 100 can be thought of as a bundle of ten	
		tens — called a "hundred."	
		b. The numbers 100, 200, 300, 400, 500,	
		600, 700, 800, 900 refer to one, two,	
		three, four, five, six, seven, eight, or nine	
		hundreds (and 0 tens and 0 ones).	
		2.NBT.B.7	
		Add and subtract within 1000, using concrete	
		models or drawings and strategies based on	
		place value, properties of operations, and/or	
		the relationship between addition and	
		subtraction; justify the reasoning used with a	
		written explanation. Understand that in	
		adding or subtracting three- digit numbers,	
		one adds or subtracts hundreds and	
		hundreds, tens and tens, ones and ones; and	
		sometimes it is necessary to compose or	
		decompose tens or hundreds.	
2.NBT.B.7		2.NBT.A.1	2.NBT.B.8
Add and subtract within 1000, using concrete		Understand that the three digits of a three-	Mentally add 10 or 100 to a given number 100-
models or drawings and strategies based on		digit number represent amounts of hundreds,	900, and mentally subtract 10 or 100 from a
place value, properties of operations, and/or		tens, and ones; e.g., 706 equals 7 hundreds, 0	given number 100–900.
the relationship between addition and		tens, and 6 ones. Understand the following as	2.NBT.B.9
subtraction; justify the reasoning used with a		special cases:	Explain why addition and subtraction strategies
written explanation. Understand that in		a. 100 can be thought of as a bundle of ten	work, using place value and the properties of
adding or subtracting three- digit numbers,		tens — called a "hundred."	operations.
one adds or subtracts hundreds and		b. The numbers 100, 200, 300, 400, 500,	
hundreds, tens and tens, ones and ones; and		600, 700, 800, 900 refer to one, two,	
sometimes it is necessary to compose or		three, four, five, six, seven, eight, or nine	
decompose tens or hundreds.		hundreds (and 0 tens and 0 ones).	

2 <sup>nd</sup> Grade Standard	Previous Grade(s) Standards	2 <sup>nd</sup> Grade Standards Taught in Advance	2 <sup>nd</sup> Grade Standards Taught Concurrently
2.NBT.B.8 Mentally add 10 or 100 to a given number 100–900, and mentally subtract 10 or 100 from a given number 100–900.		<ul> <li>2.NBT.A.1 Understand that the three digits of a three- digit number represent amounts of hundreds, tens, and ones; e.g., 706 equals 7 hundreds, 0 tens, and 6 ones. Understand the following as special cases:</li> <li>a. 100 can be thought of as a bundle of ten tens — called a "hundred."</li> <li>b. The numbers 100, 200, 300, 400, 500, 600, 700, 800, 900 refer to one, two, three, four, five, six, seven, eight, or nine hundreds (and 0 tens and 0 ones).</li> </ul>	2.NBT.B.7 Add and subtract within 1000, using concrete models or drawings and strategies based on place value, properties of operations, and/or the relationship between addition and subtraction; justify the reasoning used with a written explanation. Understand that in adding or subtracting three- digit numbers, one adds or subtracts hundreds and hundreds, tens and tens, ones and ones; and sometimes it is necessary to compose or decompose tens or hundreds. 2.NBT.B.9 Explain why addition and subtraction strategies work, using place value and the properties of operations.
2.NBT.B.9 Explain why addition and subtraction strategies work, using place value and the properties of operations.	<b>1.OA.B.3</b> Apply properties of operations to add and subtract. Examples: If $8 + 3 = 11$ is known, then $3 + 8 = 11$ is also known. (Commutative property of addition.) To add $2 + 6 + 4$ , the second two numbers can be added to make a ten, so $2 + 6 + 4 = 2 + 10 = 12$ . (Associative property of addition.) <b>1.OA.B.4</b> Understand subtraction as an unknown-addend problem. For example, subtract $10 - 8$ by finding the number that makes $10$ when added to $8$ .	<ul> <li>2.NBT.A.1</li> <li>Understand that the three digits of a three- digit number represent amounts of hundreds, tens, and ones; e.g., 706 equals 7 hundreds, 0 tens, and 6 ones. Understand the following as special cases:</li> <li>a. 100 can be thought of as a bundle of ten tens — called a "hundred."</li> <li>b. The numbers 100, 200, 300, 400, 500, 600, 700, 800, 900 refer to one, two, three, four, five, six, seven, eight, or nine hundreds (and 0 tens and 0 ones).</li> </ul>	2.NBT.B.7 Add and subtract within 1000, using concrete models or drawings and strategies based on place value, properties of operations, and/or the relationship between addition and subtraction; justify the reasoning used with a written explanation. Understand that in adding or subtracting three- digit numbers, one adds or subtracts hundreds and hundreds, tens and tens, ones and ones; and sometimes it is necessary to compose or decompose tens or hundreds. 2.NBT.B.8 Mentally add 10 or 100 to a given number 100– 900, and mentally subtract 10 or 100 from a given number 100–900.
<b>2.MD.A.1</b> Measure the length of an object by selecting and using appropriate tools such as rulers, yardsticks, meter sticks, and measuring tapes.	<b>1.MD.A.2</b> Express the length of an object as a whole number of length units, by laying multiple copies of a shorter object (the length unit) end to end; understand that the length measurement of an object is the number of same-size length units that span it with no gaps or overlaps. Limit to contexts where the object being measured is spanned by a whole number of length units with no gaps or overlaps.		

2 <sup>nd</sup> Grade Standard	Previous Grade(s) Standards	2 <sup>nd</sup> Grade Standards Taught in Advance	2 <sup>nd</sup> Grade Standards Taught Concurrently
2.MD.A.2 Measure the length of an object twice, using length units of different lengths for the two measurements; describe how the two measurements relate to the size of the unit chosen.		2.MD.A.1 Measure the length of an object by selecting and using appropriate tools such as rulers, yardsticks, meter sticks, and measuring tapes. 2.MD.A.3 Estimate lengths using units of inches, feet, centimeters, and meters.	
2.MD.A.3 Estimate lengths using units of inches, feet, centimeters, and meters.		2.MD.A.1 Measure the length of an object by selecting and using appropriate tools such as rulers, yardsticks, meter sticks, and measuring tapes.	
2.MD.A.4 Measure to determine how much longer one object is than another, expressing the length difference in terms of a standard length unit.		2.MD.A.3 Estimate lengths using units of inches, feet, centimeters, and meters.	
2.MD.B.5 Use addition and subtraction within 100 to solve word problems involving lengths that are given in the same units, e.g., by using drawings (such as drawings of rulers) and equations with a symbol for the unknown number to represent the problem.		2.MD.A.4 Measure to determine how much longer one object is than another, expressing the length difference in terms of a standard length unit.	2.OA.A.1 Use addition and subtraction within 100 to solve one- and two-step word problems involving situations of adding to, taking from, putting together, taking apart, and comparing, with unknowns in all positions, e.g., by using drawings and equations with a symbol for the unknown number to represent the problem. 2.MD.B.6 Represent whole numbers as lengths from 0 on a number line diagram with equally spaced points corresponding to the numbers 0, 1, 2,, and represent whole-number sums and differences within 100 on a number line diagram.
2.MD.B.6 Represent whole numbers as lengths from 0 on a number line diagram with equally spaced points corresponding to the numbers 0, 1, 2, , and represent whole-number sums and differences within 100 on a number line diagram.			<ul> <li>2.MD.B.5</li> <li>Use addition and subtraction within 100 to solve word problems involving lengths that are given in the same units, e.g., by using drawings (such as drawings of rulers) and equations with a symbol for the unknown number to represent the problem.</li> <li>2.MD.D.9</li> <li>Generate measurement data by measuring lengths of several objects to the nearest whole unit, or by making repeated measurements of the same object. Show the measurements by making a line plot, where the horizontal scale is marked off in whole-number units.</li> </ul>

2 <sup>nd</sup> Grade Standard	Previous Grade(s) Standards	2 <sup>nd</sup> Grade Standards Taught in Advance	2 <sup>nd</sup> Grade Standards Taught Concurrently
2.MD.C.7 Tell and write time from analog and digital clocks to the nearest five minutes, using a.m. and p.m.	1.MD.B.3 Tell and write time in hours and half-hours using analog and digital clocks.		
2.MD.C.8 Solve word problems involving dollar bills, quarters, dimes, nickels, and pennies, using \$ and ¢ symbols appropriately. <i>Example: If you</i> <i>have 2 dimes and 3 pennies, how many cents</i> <i>do you have?</i>			2.OA.A.1 Use addition and subtraction within 100 to solve one- and two-step word problems involving situations of adding to, taking from, putting together, taking apart, and comparing, with unknowns in all positions, e.g., by using drawings and equations with a symbol for the unknown number to represent the problem.
<b>2.MD.D.9</b> Generate measurement data by measuring lengths of several objects to the nearest whole unit, or by making repeated measurements of the same object. Show the measurements by making a line plot, where the horizontal scale is marked off in whole- number units.			<b>2.MD.B.6</b> Represent whole numbers as lengths from 0 on a number line diagram with equally spaced points corresponding to the numbers 0, 1, 2,, and represent whole-number sums and differences within 100 on a number line diagram.
<b>2.MD.D.10</b> Draw a picture graph and a bar graph (with single-unit scale) to represent a data set with up to four categories. Solve simple puttogether, take-apart, and compare problems using information presented in a bar graph.	<b>1.MD.C.4</b> Organize, represent, and interpret data with up to three categories; ask and answer questions about the total number of data points, how many in each category, and how many more or less are in one category than in another.		2.OA.A.1 Use addition and subtraction within 100 to solve one- and two-step word problems involving situations of adding to, taking from, putting together, taking apart, and comparing, with unknowns in all positions, e.g., by using drawings and equations with a symbol for the unknown number to represent the problem.
2.G.A.1 Recognize and draw shapes having specified attributes, such as a given number of angles or a given number of equal faces. Identify triangles, quadrilaterals, pentagons, hexagons, and cubes.	<b>1.G.A.1</b> Distinguish between defining attributes (e.g., triangles are closed and three-sided) versus non-defining attributes (e.g., color, orientation, overall size); build and draw shapes to possess defining attributes.		
2.G.A.2 Partition a rectangle into rows and columns of same-size squares and count to find the total number of them.			
2.G.A.3 Partition circles and rectangles into two, three, or four equal shares, describe the shares using the words halves, thirds, half of, a third of, etc., and describe the whole as two halves, three thirds, four fourths. Recognize that equal shares of identical wholes need not have the same shape.	<b>1.G.A.3</b> Partition circles and rectangles into two and four equal shares, describe the shares using the words halves, fourths, and quarters, and use the phrases half of, fourth of, and quarter of. Describe the whole as two of, or four of the shares. Understand for these examples that decomposing into more equal shares creates smaller shares.	2.G.A.2 Partition a rectangle into rows and columns of same-size squares and count to find the total number of them.	